

Chandra Observation of the Unidentified TeV Gamma-Ray Source HESS J1303–631 in the Galactic Plane

R. Mukherjee¹ & J. P. Halpern²

ABSTRACT

The imaging atmospheric Cherenkov array H.E.S.S. recently discovered an extended source in the 0.4–10 TeV energy range, HESS J1303–631. We obtained a 5 ks observation with the ACIS-I array on the *Chandra* X-ray observatory that does not reveal an obvious compact or diffuse X-ray counterpart. Archival *ROSAT* images are also blank in this region. Although there are several radio pulsars within the field of HESS J1303–631, none is detected in X-rays to a flux limit of $< 5 \times 10^{-14}$ ergs cm⁻² s⁻¹, and none is a likely counterpart on energetic grounds. Over the entire $17' \times 17'$ ACIS-I field, we place an upper limit of $< 5.4 \times 10^{-12}$ ergs cm⁻² s⁻¹ on the excess diffuse flux in the 2–10 keV band. One hard point-source with flux $\approx 4 \times 10^{-14}$ ergs cm⁻² s⁻¹ lies within $0'.5$ of the centroid of the TeV emission. These exploratory observations suggest that deeper pointings with *Chandra* and *XMM-Newton* are needed before we can learn more about the nature of HESS J1303–631. Its similarity to the unidentified source TeV J2032+4130 indicates the probable existence of a new class of high-energy source in the Galactic plane that originates from young, massive stars or their supernova remnants.

Subject headings: gamma-rays: individual (HESS J1303–631) — gamma-rays: observations — X-rays: stars

1. Introduction

The new imaging atmospheric Cherenkov telescope (IACT) H.E.S.S. (High Energy Stereoscopic System) is located in Namibia and is sensitive in the hundreds of GeV regime (Hinton 2004). The complete H.E.S.S. array of four telescopes has been operational since

¹Department of Physics & Astronomy, Barnard College, Columbia University, New York, NY 10027

²Department of Astronomy, Columbia University, New York, NY 10027

2003 December and has reported observations of several astrophysical sources (e.g., Aharonian et al. 2004; Beilicke et al. 2004a). Recent results include the discovery of a new, unidentified TeV γ -ray source HESS J1303–631 (Beilicke et al. 2004a; Aharonian et al. 2005a). This is only the second unidentified source at TeV energies in the Galactic plane, at $(l, b) = (304.^{\circ}24, -0.^{\circ}36)$. HESS J1303–631 was discovered in the pointings at the Be/X-ray pulsar PSR B1259–63 (Beilicke et al. 2004b) for 39 hr of total exposure from 2004 February to June. The new source is located $0.^{\circ}6$ north of the pulsar, at R.A. = $13^{\text{h}}03^{\text{m}}00.^{\text{s}}4 \pm 04.^{\text{s}}4$, decl. = $-63^{\circ}11'55'' \pm 31''$ (J2000.0), and was observed to be brighter than the pulsar, about 0.17 crab, or 1.2×10^{-11} ergs cm^{-2} s^{-1} above 380 GeV. The statistical significance of detection was 21σ . There is good evidence that the TeV emission is extended, with an intrinsic Gaussian $\sigma = 0.^{\circ}16 \pm 0.^{\circ}02$. There is no evidence of flux variability among the H.E.S.S. observations. These properties are similar to those of TeV J2032+4130, the first such unidentified source that was discovered by the HEGRA CT system (Aharonian et al. 2002). Also in the Galactic plane at $(l, b) = (80.^{\circ}25, +1.^{\circ}07)$, TeV J2032+4130 lies in the direction of the Cygnus OB2 stellar association, and its nature remains uncertain despite extensive multiwavelength efforts to find a counterpart (Mukherjee et al. 2003; Butt et al. 2003).

In this paper we report on initial efforts to identify HESS J1303–631. Multiwavelength searches of individual EGRET source fields were often successful in finding a likely identification (see Mukherjee & Halpern 2004, for a review). A difficulty faced by the EGRET searches, especially in the Galactic plane, is the relatively large positional uncertainty, $\approx 0.^{\circ}5 - 1.^{\circ}$. IACTs like H.E.S.S. or HEGRA have the advantage of much better angular resolution resulting in smaller error boxes for γ -ray source positions in comparison to EGRET, even when the extent of HESS J1303–631 is included.

2. High-Energy and Radio Observations

2.1. *ROSAT*

Prior to the *Chandra* observation reported here, the error box of HESS J1303–631 was covered only by a *ROSAT* observation, which does not show any bright sources in the region of interest. No pointings by *ASCA* or *BeppoSAX* were made in this region. Since this source is in the Galactic plane (similar to TeV J2032+4130), it is likely that its X-ray counterpart is highly absorbed. The total H I column density in this direction is $N_{\text{H}} = 2 \times 10^{22}$ cm^{-2} , which, not even including molecular hydrogen, has optical depth > 1 at $E < 3$ keV, and should be associated with visual extinction $A_V \approx 11$ mag (Predehl & Schmitt 1995). Figure 1 shows the X-ray image taken with the *ROSAT* Position Sensitive Proportional Counter (PSPC) in the energy range 0.2–2.0 keV, covering the field of HESS J1303–631. This image

was created by co-adding exposure-corrected skymaps of 49 ks of data. The original target of the *ROSAT* observations was PSR B1259–63. The small circle corresponds to the 1σ uncertainty of the centroid of HESS J1303–631, and includes both statistical errors and pointing (systematic) errors. The systematic pointing uncertainty of the H.E.S.S. telescopes is estimated to be $\sim 20''$ (Aharonian et al. 2005a). The statistical and systematic errors were added in quadrature. The large circle around the TeV position indicates the Gaussian $\sigma = 0.16$ extent of the TeV source (Aharonian et al. 2005a). The only significant X-ray source in Figure 1 is the K0 star HD 113182 at R.A. = $13^{\text{h}}02^{\text{m}}59.^{\text{s}}3$, decl. = $-63^{\circ}27'44''$ (J2000.0), which is inconsistent with the H.E.S.S. location. The numbered triangles in the region of the H.E.S.S. source are the locations of radio pulsars from the Parkes Multibeam Survey. Their properties are given in Table 1. While there are several radio pulsars near the H.E.S.S. position, none is detected in the *ROSAT* image. A weak, out-of-focus *ROSAT* source near the location of PSR J1301–6310 ($d \approx 2.1$ kpc) is detected only in soft X-rays, indicating that it is probably from a foreground star, and not attributable to the pulsar. Within the Gaussian σ extent of the H.E.S.S. source, there are no other candidate radio sources in published surveys (Aharonian et al. 2005a).

2.2. *Chandra*

We observed the location of HESS J1303–631 on 2004 September 25 with the Advanced CCD Imaging Spectrometer (ACIS, Burke et al. 1997) onboard the *Chandra* X-ray Observatory (Weisskopf, O’Dell, & van Speybroeck 1996; Weisskopf et al. 2002) using Director’s discretionary time. A preliminary estimate of the source centroid was positioned at the default location on the front-illuminated ACIS-I array. The standard timed readout with a frame time of 3.2 s was used, and the data were collected in VFaint mode. A total of 4707 s of on-time was accumulated, while the effective exposure live-time was 4648 s. After processing the event file through the VFaint mode filter, a total of 4188 diffuse events in the 2 – 10 keV band were found on the four ACIS-I CCDs in addition to five point sources with seven or more photons each. Figure 2 shows the *Chandra* image with the five point sources marked. Their positions and count rates are listed in Table 2, together with possible stellar optical counterparts. None of the radio pulsars listed in Table 1 was detected in the *Chandra* observation, to a limit of $\approx 5 \times 10^{-14}$ ergs cm $^{-2}$ s $^{-1}$ in the 2 – 10 keV band.

The one hard X-ray source (#7) without a possible optical (or 2MASS) counterpart lies only $32''$ from the centroid of the TeV source, which is consistent with the combined $20''$ systematic and $31''$ statistical uncertainties of the latter. Assuming a power-law spectrum of photon index $\Gamma = 2.0$ and column density $N_{\text{H}} = 2 \times 10^{22}$ cm $^{-2}$ (the total H I column density

in this direction), source 7 has a $2 - 10$ keV flux of $\approx 4 \times 10^{-14}$ ergs cm $^{-2}$ s $^{-1}$. It could be a background AGN, or a candidate for identification with HESS J1303–631.

Any fainter point sources than the ones listed in Table 2 are of marginal significance, and are included in the following analysis of the diffuse flux. Although no extended source is apparent in the *Chandra* image, we use the standard quiescent background rates to estimate that 3239 of the diffuse events in the $2 - 10$ keV band are particle plus cosmic X-ray background, leaving 949 diffuse photons over the $17' \times 17'$ ACIS-I field. These 949 photons may be Galactic diffuse background or diffuse flux originating from HESS J1303–631. Conservatively, this 0.21 photons s $^{-1}$ of diffuse flux can be considered an upper limit to the flux of HESS J1303–631. Assuming a power-law spectrum of photon index $\Gamma = 2.0$ and column density $N_{\text{H}} = 2 \times 10^{22}$ cm $^{-2}$, this corresponds to a $2 - 10$ keV observed flux of $< 5.4 \times 10^{-12}$ ergs cm $^{-2}$ s $^{-1}$, or an unabsorbed $2 - 10$ keV flux of $< 6.4 \times 10^{-12}$ ergs cm $^{-2}$ s $^{-1}$.

If we perform the same analysis separately in the $2 - 5$ keV band, where the instrumental background is lower and the subtraction less subject to systematic effects, we find an observed diffuse flux of $< 2.2 \times 10^{-12}$ ergs cm $^{-2}$ s $^{-1}$, or an unabsorbed $2 - 5$ keV flux of $< 3.0 \times 10^{-12}$ ergs cm $^{-2}$ s $^{-1}$. These limits would be reduced by a factor of several if the diffuse source is considerably smaller than the $17' \times 17'$ ACIS-I field. The fluxes limits are subject to additional uncertainty from the unknown shape of the X-ray spectrum, and possible column density of intervening molecular material would increase the limits.

2.3. EGRET

HESS J1303–631 was not detected by EGRET above 100 MeV. The source is located in the Galactic plane, which has intense diffuse gamma-ray emission. For the Galactic plane sources, the third EGRET (3EG) Catalog (Hartman et al. 1999) lists point sources that were detected at least 5σ above the diffuse model. We have estimated the upper limit of γ -ray emission to be 10.4×10^{-8} photons cm $^{-2}$ s $^{-1}$ above 100 MeV.

3. Discussion

The short *Chandra* observation presented here does not reveal much about the nature of HESS J1303–631. Those point sources detected primarily in soft X-rays (< 2 keV) are likely to be coronal, with late-type stellar counterparts identified in Table 2, and unrelated to HESS J1303–631. The one hard X-ray source without an optical counterpart (#7 in Table 2) could be a background AGN. The position of source 7 is consistent with the well-

determined (0'.5) location of the centroid of the TeV emission. However, the extended and steady nature of HESS J1303–631 disfavors a blazar origin. The absence of a brighter X-ray counterpart argues against a microquasar origin, but perhaps allows an isolated compact object.

No diffuse source is apparent in the *Chandra* image, thus, there is no further evidence concerning the extended TeV emission, and whether it has an electronic or hadronic origin. Based on their spin-down fluxes $\dot{E}/4\pi d^2$, none of the known pulsars in the HESS J1303–631 field (Table 1) is a good candidate for powering detectable TeV emission from a pulsar wind nebula (PWN). The highest spin-down flux is from PSR J1301–6305, which has $\dot{E}/4\pi d^2 = 5.6 \times 10^{-11}$ ergs cm⁻² s⁻¹, and would imply an unreasonably high efficiency of $\approx 38\%$ for production of 0.3 – 10 TeV flux, which is 2.1×10^{-11} ergs cm⁻² s⁻¹ according to the extrapolated spectrum (Aharonian et al. 2005a). In comparison, *Chandra* sets an upper limit on the 2–10 keV flux of each of these pulsars of $\approx 5 \times 10^{-14}$ ergs cm⁻² s⁻¹, which is consistent with the normal range of $10^{-4} \leq L_x/\dot{E} \leq 10^{-2}$ for pulsars (Possenti et al. 2002). Furthermore, there is an empirical threshold $\dot{E}$ of $\approx 4 \times 10^{36}$ ergs s⁻¹ below which pulsars do not power bright PWN either in X-rays or in radio (Gotthelf 2004). None of the pulsars in Table 1, including PSR J1301–6305, exceeds this threshold.

It is instructive to compare this *Chandra* observation of HESS J1303–631 to a similar 5 ks *Chandra* observation of the HEGRA source TeV J2032+4130, the first unidentified TeV source. TeV J2032+4130 is located near the Cygnus OB2 region, in which several of the stars are among the most active stellar X-ray sources in the Galaxy. In contrast to HESS J1303–631, the TeV J2032+4130 field contains more point X-ray sources, spectroscopic optical identifications of which indicated that they are mostly a mix of O-type stars in the Cyg OB2 association, at $d \approx 1.7$ kpc, and foreground late-type stars (Mukherjee et al. 2003). The only unusual X-ray source in the field of TeV J2032+4130 is a highly variable, hard and absorbed X-ray source, the brightest in the *Chandra* image, located 7' from the centroid of TeV J2032+4130. It is not clear if this unidentified X-ray source is associated with TeV J2032+4130. If so, Mukherjee et al. (2003) argued that it would be located considerably beyond Cyg OB2, and still be within the young Galactic disk.

It has been argued that colliding stellar winds in the Cyg OB2 association could be responsible for the production of TeV gamma-rays in TeV J2032+4130 (Bengalia et al. 2001). Aharonian et al. (2002) hypothesized two possible origins for extended TeV emission that may be displaced from its originating source of energy. One is π^0 decay resulting from hadrons accelerated in shocked OB star winds and interacting with a local, dense cloud. The other is TeV emission in a jet-driven termination shock due to Inverse Compton scattering, either from an as-yet undetected microquasar, or from Cyg X-3. Torres, Domingo-Santamaria, &

Romero (2004) theorized that groups of stars in OB associations located close to cosmic ray acceleration sites may be sources of TeV emission, and cited TeV J2032+4130 as a candidate for such a source. In such a scenario, no MeV-GeV emission is expected from the source at EGRET or *GLAST* energies.

The HEGRA source TeV J2032+4130 also showed some evidence for spatial extent, with a Gaussian 1σ radius of $5'.6 \pm 1'.7$ (Aharonian et al. 2002). As in the case of HESS J1303–631, the HEGRA source was found to be steady in observations from 1999 to 2001, unlike the flaring blazars detected by EGRET and the IACTs. However, unlike TeV J2032+4130, HESS J1303–631 is not located near a dense OB stellar association or a star formation region. Although Aharonian et al. (2005a) noted that HESS J1303–631 is superposed on the Centaurus OB1 association, this is a rather sparse group of only 20 stars, mostly of type B, spread over $\approx 6^\circ$ or 260 pc at the adopted distance of 2.5 kpc (Humphreys 1978) (2.3 kpc according to Kaltcheva & Georgiev 1994). The nearest member is 0.8° from HESS J1303–631. Therefore, we hesitate to posit a physical origin of the small TeV source from the group as a whole, or even from any existing member. Also, the average visual extinction to stars in Cen OB1 is only $A_V \approx 2.0$ mag (Kaltcheva & Georgiev 1994), which is not large enough to account for the absence of an X-ray counterpart at this distance. However, it is possible that a deceased member of the association, whose compact remnant is difficult to detect, is responsible for HESS J1303–631.

4. Conclusions

That both TeV J2032+4130 and HESS J1303–631 are indeed Galactic now seems secure. It is likely that several more such sources will be detected in future sky surveys to be conducted by H.E.S.S. and VERITAS. Such TeV sources may not be associated with the numerous unidentified EGRET sources in the 3EG catalog, and could very well comprise a new class by themselves. Given the apparent latitude range of $\sim 1^\circ$, their scale height must be < 200 pc if located at distances < 10 kpc. Thus, they are not very old, and probably originate from massive stars or their remnants.

After this paper was submitted, the discovery by H.E.S.S. of eight additional sources in a survey of the Galactic plane was reported (Aharonian et al. 2005b). Their mean Galactic latitude of -0.25° and rms dispersion in latitude of 0.25° further supports an association with young, massive stars. All of these new sources appear to be extended, and some are plausibly associated with known supernova remnants, which suggests that the remaining unidentified sources might also be supernova remnants that have not yet been detected or recognized at longer wavelengths because of their considerable distances and/or weak synchrotron emission.

The search for and evaluation of candidate counterparts will be very important in resolving the nature of these TeV sources.

We thank Harvey Tananbaum for granting the *Chandra* Director’s discretionary time for this observation. This work was supported in part by the National Science Foundation.

REFERENCES

- Aharonian, F., et al. 2002, A&A, 393, 37L
- . 2004, A&A, 425, L13
- . 2005a, A&A, submitted
- . 2005b, Science, 307, 1938
- Beilicke, M., Khelifi, B., Masterson, C., de Naurois M., & Schlenker, S. 2004a, presented at the Int. Symp. on High Energy Gamma-ray Astronomy, Heidelberg (July 2004); F. A. Aharonian, H. J. Völk, D. Horns, Eds. (AIP Conference Proceedings 745, 2005), pg. 347
- Beilicke, M., Ouchrif, M., Rowell, G., & Schlenker, S. 2004b, IAU Circ. 8300, 2
- Bengalia, P., Romero, G. E., Stevens, I. R., & Torres, D. F. 2001, A&A, 366, 605
- Burke, B. E., Gregory, J., Bautz, M. W., Prigozhin, G. Y., Kissel, S. E., Kosicki, B. N., Loomis, A. H., & Young, D. J. 1997, IEEE Trans. Electron Devices, 44, 1633
- Butt, Y. M., et al. 2003, ApJ, 597, 494
- Gotthelf, E. V. 2004, in Young Neutron Stars and their Environments, IAU Symp. 218, ed. F. Camilo & B. M. Gaensler (San Francisco: ASP), 225
- Hartman, R. C., et al. 1999, ApJS, 123, 79
- Hinton, J. A. 2004, New Astron. Rev., 48, 331
- Humphreys, R. M. 1978, ApJS, 38, 309
- Kaltcheva, N. T., & Georgiev, L. N. 1994, MNRAS, 269, 289
- Kramer, M., et al. 2003, MNRAS, 342, 1299

- Manchester, R. N., et al. 2001, MNRAS, 328, 17
- Mukherjee, R., & Halpern, J. 2004, in “Cosmic Gamma-ray Sources”, ed. K.S. Cheng & G.E. Romero (Dordrecht: Kluwer), in press (astro-ph/0408063)
- Mukherjee, R., Halpern, J. P., Gotthelf, E. V., Eracleous, M., & Mirabal, N. 2003, ApJ, 589, 487
- Possenti, A., Cerutti, R., Colpi, M., Merghetti, S. 2002, A&A, 387, 993
- Predehl, P., & Schmitt, J. H. M. M. 1995, A&A, 293, 889
- Torres, D. F., Domingo-Santamaria, E., & Romero, G. E. 2004, ApJ, 601, 75L
- Weisskopf, M. C., O’Dell, S. L., & van Speybroeck, L. P. 1996, Proc. SPIE, 2805, 2
- Weisskopf, M. C., Brinkman, B., Canizares, C., Garmire, G., Murray, S., & van Speybroeck, L. P. 2002, PASP, 114, 1

Table 1. Radio Pulsars in the Field of HESS J1303–631

Ident.	Name	Position		P (s)	$\log \tau$ (yr)	$\log \dot{E}$ (ergs s ⁻¹)	d (kpc)	$\log(\dot{E}/4\pi d^2)$ (ergs cm ⁻² s ⁻¹)	Ref
		R.A.	Decl.						
1	PSR J1301–6310	13 01 28.3	–63 10 40	0.664	5.27	33.88	2.1	–10.83	1
2	PSR J1301–6305	13 01 45.8	–63 05 34	0.185	4.04	36.23	15.8	–10.25	2
3	PSR J1302–6313	13 02 19.2	–63 13 29	0.968	6.38	32.44	28.1	–14.53	1
4	PSR J1303–6305	13 03 00.0	–63 05 01	2.307	7.22	30.85	13.6	–15.50	2

Note. — Units of right ascension are hours, minutes, and seconds. Units of declination are degrees, arcminutes, and arcseconds.

References. — (1) Kramer et al. 2003; (2) Manchester et al. 2001.

Table 2. *Chandra* Sources in the Field of HESS J1303–631

<i>Chandra</i> Ident.	X-ray Position		Counts (< 2 keV)	Counts (> 2 keV)	Nearest USNO B1.0 Star		<i>B</i> (mag)	<i>R</i> (mag)	Dist ($''$)
	R.A.	Decl.			R.A.	Decl.			
5	13 02 09.67	–63 08 08.4	9	0	13 02 09.53	–63 08 09.1	18.40	16.02	1.2
6	13 02 26.35	–63 05 43.0	9	0	13 02 26.42	–63 05 42.4	18.59	16.10	0.8
7	13 03 00.98	–63 11 23.2	0	7	...	...	...	...	...
8	13 03 38.59	–63 16 58.5	7	0	13 03 38.71	–63 16 59.2	15.01	14.32	1.1
9	13 03 41.93	–63 09 13.5	13	5	13 03 41.92	–63 09 12.9	19.68	16.52	0.6
					13 03 42.11	–63 09 13.8	16.46	15.78	1.2

Note. — Units of right ascension are hours, minutes, and seconds. Units of declination are degrees, arcminutes, and arcseconds.




Fig. 1.— *ROSAT* PSPC X-ray image of the field of HESS J1303–631. The small circle corresponds to the 1σ uncertainty of the centroid of the H.E.S.S. source, which includes both statistical errors and pointing (systematic) errors. The large circle is the estimated Gaussian 1σ extent of the TeV emission (Aharonian et al. 2005a). The squares are the fields of view of the CCDs in the subsequent *Chandra* observation (see Figure 2). The numbered triangles are the locations of radio pulsars from the Parkes Multibeam Survey, whose properties are listed in Table 1.




Fig. 2.— *Chandra* ACIS-I image of the field of HESS J1303–631. The numbered triangles mark the locations of radio pulsars from the Parkes Multibeam Survey, whose properties are listed in Table 1. The small circle corresponds to the 1σ uncertainty of the centroid of HESS J1303–631, and includes both statistical errors and pointing (systematic) errors. The large circle is the estimated Gaussian 1σ extent of the TeV emission (Aharonian et al. 2005a). The plus signs mark the significant *Chandra* point sources, whose properties are listed in Table 2. Source 7 is $32''$ from the nominal centroid of the TeV emission.